



**West African Ornithological Society
Société d'Ornithologie de l'Ouest
Africain**



**Join the WAOS and support
the future availability of free
pdfs on this website.**

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.
If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

**Devenez membre de la
SOOA et soutenez la
disponibilité future des pdfs
gratuits sur ce site.**

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.
Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante
(2) pour économiser du papier.

ADDITIONS TO LOCAL AVIFAUNAS: ZARIA - The following birds new to the Zaria area have been reported since the additions made by Dyer (*Bull. Niger. Orn. Soc.* 13, 1977, 82-83). Observers: M.D. - M. Dyer, M.E.G. - M. E. Gartshore, A.U.E. - A. U. Ezealor, C.R. - C. Ryall, H.Q.P.C. - H. Q. P. Crick.

Glossy Ibis *Plegadis falcinellus* About 20 over University Farm Lake in early April, 1981 (M.D., M.E.G.).

Wigeon *Anas penelope* A female feeding with Garganey *A. querquedula* at University Farm Lake, 10-12 April 1981 (M.D., M.E.G., H.F.C.).

Booted Eagle *Hieraaetus pennatus* One over experimental plots near Bomo Lake, Samaru, Zaria, on 31 January 1981 (M.D., M.E.G., A.U.E.).

Long-legged Buzzard *Buteo rufinus* Observed at the same time as the above, this distinctive raptor circled over fields and alighted, allowing very close approach (M.D., M.E.G., A.U.E.).

Scaly Francolin *Francolinus squamatus* A pair heard duetting at about 0800 hrs., 5 August 1981 in dense savanna woodland adjacent to gallery forest at Anara For. Res. 5 August 1981. This observation represents a substantial range extension north in Nigerian savannas. (It is common at Nindam For. Res. near Kafanchan and Kagoro (M.E.G., C.R.)).

Finfoot *Podica senegalensis* An immature bird was purchased from fisherman at Kubanni reservoir in February 1979. It was later released at University Farm Lake (M.D., M.E.G.).

Kittlitz's Sandplover *Charadrius pecuarius* Six feeding on shore of Kubanni reservoir in early April 1981 (M.D.).

Avocet *Recurvirostra avosetta* Two in winter plumage on Kubanni reservoir, late March 1981 (M.D., M.E.G.).

Black-headed Gull *Larus ridibundus* Its inland status in West Africa has changed in recent years (Sharland Malimbus 3, 1981, 54) and it is now a frequent visitor to Zaria reservoirs in the dry season (M.E.G., M.D.).

Black Tern *Sterna nigra* One with a White-winged Tern *S. leucoptera* at Samaru sewage works in early April 1981 (distinguished by marginally smaller size, darker colouration, dark pectoral area and greyish tail) (M.E.G.).

Yellowbill *Ceuthmochanes aereus* Observed on several occasions in gallery forest at Anara Forest Reserve; also at Old Birnin Gwarri (just outside the Zaria area) (M.D., M.E.G.).

Narina Trogon *Apaloderma narina* At least three observed throughout the wet season in dense gallery forest at Anara Forest Reserve (M.E.G., A.U.E., C.R.).

Yellow-bellied Bunting *Emberiza flaviventris* A pair in August 1981, the male singing (M.E.G.).

Melba Finch *Pytilia melba* Seen on several occasions from August, 1979 to August 1981 at Samaru also in June 1980. This species is normally associated with Sudan savanna and may be shifting southward in response to

changes wrought by more intensive cultivation (M.D., M.E.G.).

Black-bellied See cracker *Pirenestes ostrinus* Reported by Hartert (Novit. Zool. 22, 1915, 244-266) as a breeding resident near Kaduna, its presence at Zaria has never been substantiated. Several were observed in tall grass in a fadama area at Anara Forest Reserve in September 1980 (M.D., H.Q.P.C.).

M. E. Gartshore

606 Martins Road, RR3, Dundas, Ontario L9H 5E3

DESTRUCTION OF EUROPEAN WHITE STORKS IN NIGERIA BY SHOOTING - Even more noteworthy that the number of Spanish-ringed White Storks *Ciconia ciconia* recovered in Nigeria in the last two years - eight - is their fate there. One was caught and released alive with its original ring; the other seven were shot. Details are:

C 15540 Ringed 07/06/1977 (pullus), Miajadas, Cáceres, Spain 39°09'N 05°54'W
-/12/1890, Nguru, Nigeria 12°53'N 10°30'E Shot

C 14334 26/05/1978 (pullus), El Gordo, Cáceres, 39°52'N 05°20'W
-/01/1980, Gashua, Nigeria 12°53'N 11°05'E Released alive

C 11390 26/05/1978 (pullus), Elmaraz, Spain 39°50'N 05°37'W
-/12/1980, Nguru Shot

C 18943 27/05/1978 (pullus), El Gordo
19/01/1980 Kano 12°55'N 10°30'E Nigeria Shot

C 18823 05/06/1979 (pullus) Brozas, Cáceres 39°37'N 06°47'W
-/12/1981, Nguru Shot

G 01351 06/06/1980 (pullus), Muñana, Avila, Spain 40°35'N 05°03'W
-/12/1981, Nguru Shot

G 03338 04/06/1981 (pullus), Casas Viejas, Cadiz, Spain 36°21'N 05°49'W
-/12/1981, Nguru Shot

G 03491 06/06/1981 (pullus), Casas Viejas, Cadiz
-/12/1981, Nguru Shot

All of the indications are that the Iberian/Maghreb population of White Storks, separated from the larger east European population by an ever-increasing distance, is in a state of decline (S. Cramp *The Birds of the Western Palearctic*, 1977. See p. 329 and map p. 331). Doubtless the principal cause is drainage of European wetlands, but unnecessary persecution in African winter quarters cannot help. 17 other White Storks have been recovered in Nigeria since 1964 (R. E. Sharland, Ringing in Nigeria ... Annual Reports Nos. 9, 10, 11, 13, 14, 15-16, 17, 21 and 23, *Bull. Niger. Orn. Soc. and Malimbus*), ringed in Spain (6), Tunisia (3), Morocco (1), Algeria (1), France (2), Germany (3) and Greece (1); two of them were recovered at Azare (39°37'N 22°16'E) (one "killed" and the other "caught by trapper and later released"), one at Biu (10°36'N 12°11'E), and the remaining 14 at Nguru. The fate of the Nguru 14 has not been given but it is safe to assume that they were all or nearly all caught alive by trappers for the well-known Kano live-bird trade. Presumably that trade continues unabated (R. E. Sharland, pers. comm.), with illegal destruction by shooting a new hazard and an additional burden for the wintering White Stork population to bear.

C. H. Fry

Aberdeen University Zoology Dept., Tillydrone Ave., Aberdeen AB9 2TN, U.K.